Vaccination Report – 20 September 2022

1. Vaccine Implementation

• <u>WHO's Emergency Use Listing(EUL) Vaccines</u> (Last Updated 7 July 2022)

	Manufacturer	Name of Vaccine	NRA of Record	Vaccine type	
1	Pfizer-BioNTech (US)	BNT162b2/COMIRNATY Tozinameran (INN)	EMA,USFDA	mRNA	
2	AstraZeneca (UK)	AZD1222 Vaxzevria	EMA, MFDS KOREA, Japan MHLW/PMDA, Australia TGA, COFEPRIS(Mexico), ANMAT(Argentina)	Non ReplicatingViral vector	
3	Serum Institute of India (India)	Covishield (ChAdOx1_nCoV-19)	DCGI	Non Replicating Viral Vector	
4	Johnson &Johnson (US)	Ad26.CoV2.S	EMA, DCGI	Non ReplicatingViral vector	
5	Moderna (US)	mRNA-1273	EMA, USFDA, MFDS	mRNA	
6	Sinopharm Beijing (China)	SARS-CoV-2 Vaccine(Vero Cells)	NMPA	Inactivated virus (Vero Cells)	
7	Sinovac (China)	COVID-19 Vaccine (Vero Cells)	NMPA	Inactivated virus (Vero Cell)	
8	Bharat Biotech (India)	SARS-CoV-2 Vaccine, Inactivated (Vero Cell)/ COVAXIN	DCGI	Whole-Virion Inactivated (Vero Cell)	
9	Serum Institute of India (India)	NVX-CoV2373/Covovax	DCGI Protein Subur		
10	NÔVAVÁX (US)	NVX-CoV2373/Covovax	EMA	Protein Subunit	
11	CanSinoBIO (China)	Ad5-nCoV	NMPA	Non ReplicatingViral vector	

• **47** Vaccines Approved by at Least One Country

Vaccine Type	mRNA	Non Replicating Viral vector	Inactivated virus	Protein Subunit	DNA	Virus-like Particles (VLP)	Total
In Use	8	9	11	17	1	1	47

Source: <u>https://covid19.trackvaccines.org/vaccines/approved/#vaccine-list</u> (Last Updated 19 September 2022)


 Vaccination against COVID-19 has now started in 218 locations (Source: <u>Our World in Data</u>. Last Updated 19 September 2022)

Location	Doses Given	Complete Initial Protocol (% of population)	Partly Vaccinated (% of population)
Worldwide	12.68 billion	4.94 billion (62.46 %)	5.37 billion (67.90 %)

About this data:

a: This data changes rapidly and might not reflect doses still being reported. It may differ from other sites & sources. b: Where data for full vaccinations is available, it shows how many people have received at least 1 dose and how many people have been fully vaccinated (which may require more than 1 dose).Where data for full vaccinations isn't available, the data shows the total number of vaccine doses given to people. Since some vaccines require more than 1 dose, the number of fully vaccinated people is likely lower.

c: It only has full vaccination totals in some locations.




Note: Alternative definitions of a full vaccination, e.g. having been infected with SARS-CoV-2 and having 1 dose of a 2-dose protocol, are ignored to maximize comparability between countries.

Share of people who completed the initial COVID-19 vaccination protocol, Sep 19, 2022




Total number of people who received all doses prescribed by the initial vaccination protocol, divided by the total population of the country.



Source: Official data collated by Our World in Data – Last updated 20 September 2022 OurWorldInData.org/coronavirus • CC BY Note: Alternative definitions of a full vaccination, e.g. having been infected with SARS-CoV-2 and having 1 dose of a 2-dose protocol, are ignored to maximize comparability between countries.

COVID-19 vaccine boosters administered per 100 people, Sep 19, 2022 Total number of vaccine booster doses administered, divided by the total population of the country. Booster doses are doses administered beyond those prescribed by the original vaccination protocol.




Source: Official data collated by Our World in Data - Last updated 20 September 2022

OurWorldInData.org/coronavirus • CC BY

COVID-19 vaccination policy, Sep 19, 2022

Policies for vaccine delivery. Vulnerable groups include key workers, the clinically vulnerable, and the elderly. "Others" include select broad groups, such as by age.



Source: Oxford COVID-19 Government Response Tracker, Blavatnik School of Government, University of Oxford – Last updated 20 September 2022 OurWorldInData.org/coronavirus • CC BY

2. Effectiveness of Vaccine and/or Previous Infection against symptomatic infection for Alpha, Delta and Omicron variants

Vaccine Status	Vaccine Effectiveness			
	Alpha	Delta	Omicron	
1 Dose (BNT162b2 or ChAdOx1 nCoV-19)	48.7% (95%Cl: 45.5-51.7%) ¹ 66%(BNT162b2) ⁴ 64% (ChAdOx1) ⁴	30.7% (95%CI: 25.2-35.7%) ¹ 56% (BNT162b2) ⁴ 67% (ChAdOx1) ⁴ 82% (95% CI:73- 91%) ⁷		



BY

1 Dose (mRNA-1273)	83% ⁴	72% ⁴	
1 Dose(Sinopharm or Sinovac)	00 /0	13.8%,(95%Cl: -60.2-54.8%) ³	
2 Doses (BNT162b2)	93.7% (95%CI: 91.6-95.3) ¹ 76% (95%CI: 69-81%) ² 89% ⁴	88% (95% CI: 85.3-90.1%) ¹ 42% (95% CI: 13-62%) ² 87% ⁴ 93% (95% CI: 88-97%/12-18Y) ⁵ 93% (95% CI: 88-97%) ⁷	50% (95% CI: 35%–62%) ⁸
2 Doses (ChAdOx1 nCoV-19)	74.5% (95%CI: 68.4-79.4%) ¹	67.0% (95%CI: 61.3-71.8%) ¹	
2 Doses (mRNA-1273)	86%, (95%Cl: 81-90.6%) ²	76%, (95% Cl: 58-87%) ²	30.4% (95% CI: 5.0%-49.0%) ⁹
2 Doses(Sinopharm or Sinovac)		59.0%, (95%CI: 16.0-81.6%) ³	
3 Doses (BNT162b2)		95.33% (SD 6.44) ⁶ 86.1% (95% CI, 67.3 to 94.1) ¹¹	67.2% (95% CI: 66.5- 67.8%) at 2 to 4 weeks ¹⁰ 49.4% (95% CI, 47.1 to 51.6) ¹¹ 52.2% (95% CI, 48.1 to 55.9) ¹²
3 Doses(mRNA-1273)			62.5% (95% CI: 56.2-67.9%) ⁹ 47.3% (95% CI, 40.7 to 53.3) ¹¹
2 Doses (BNT162b2) + 1Dose(mRNA-1273)			73.9% (95% CI: 73.1-74.6%) at 2 to 4 weeks ¹⁰
2 Doses(ChAdOx1 nCoV- 19)+1Dose(BNT162b2)			62.4% (95% CI, 61.8- 63.0) at 2 to 4 weeks ¹⁰
2 Doses (ChAdOx1 nCoV-19)+ 1Dose (mRNA-1273)			70.1% (95% CI, 69.5 to 70.7) at 2 to 4 weeks ¹⁰
2 Doses (BNT162b2) +Previous infection			55.1% (95% CI, 50.9 to 58.9) ¹²
3 Doses (BNT162b2) +Previous infection			77.3% (95% CI, 72.4 to 81.4) ¹²
Previous Omicron Infection			76.1% on BA.4 or BA.5 (95% Cl: 54.9 to 87.3%) ¹³

References:

- 1) Effectiveness of Covid-19 Vaccines against the B.1.617.2 (Delta) Variant
- 2) <u>Comparison of two highly-effective mRNA vaccines for COVID-19 during periods of Alpha</u> and Delta variant prevalence
- 3) <u>Efficacy of inactivated SARS-CoV-2 vaccines against the Delta variant infection in</u> <u>Guangzhou: A test-negative case-control real-world study</u>
- 4) Effectiveness of COVID-19 vaccines against variants of concern in Ontario, Canada
- 5) Effectiveness of BNT162b2 Vaccine against Delta Variant in Adolescents
- 6) <u>A RCT of a third dose CoronaVac or BNT162b2 vaccine in adults with two doses</u> of CoronaVac
- 7) Effectiveness of BNT162b2 Vaccine against Delta Variant in Adolescents
- 8) Effectiveness of BNT162b2 Vaccine against Omicron Variant in South Africa
- 9) Effectiveness of mRNA-1273 against SARS-CoV-2 omicron and delta variants
- 10) Covid-19 Vaccine Effectiveness against the Omicron (B.1.1.529) Variant
- 11) Effect of mRNA Vaccine Boosters against SARS-CoV-2 Omicron Infection in Qatar
- 12) Effects of Previous Infection and Vaccination on Symptomatic Omicron Infections
- 13) <u>Protection of SARS-CoV-2 natural infection against reinfection with the BA.4 or BA.5</u> <u>Omicron subvariants</u>

3. Latest Relevant Articles

- <u>Effectiveness and Durability of the BNT162b2 Vaccine against Omicron</u> <u>Sublineages in South Africa (September 14, 2022)</u>
- <u>Covid-19 Vaccines Immunity, Variants, Boosters</u>(September 15, 2022)
- <u>A Bivalent Omicron-Containing Booster Vaccine against Covid-19</u> (Published September 16, 2022)
- <u>Omicron BA.2 breakthrough infection enhances cross-neutralization of BA.2.12.1</u> and BA.4/BA.5 (Published 20 September 2022)

4. Other Information

 <u>CDC</u>: Mortality Risk Among Patients Hospitalized Primarily for COVID-19 During the Omicron and Delta Variant Pandemic Periods — United States, April 2020– June 2022(Published September 16, 2022)